What should be done during the checkup?

	Newborn - 12 mo	15 mo – 4 yrs	5–10 yrs	11-21 yrs
History				
Initial/Interval	✓	✓	✓	✓
Measurements				
Height & Weight	✓	✓	✓	✓
Head circumference	✓	✓ (up to 24 mo.)		✓
Blood Pressure		✓ (start at 3 yrs)	✓	✓
Sensory Screening				
Vision	s	s	o	o
Hearing	s	s	o	s/o
Developmental / Behavioral Assessment	✓	✓	✓	✓
Immunization	✓	✓	✓	✓
Procedures-Patients at Risk				
Lead Screening	✓ (9-12 mo.)	✓ (24 mo.)	Required by age 7	
Tuberculin Test	✗ (12 mo.)	✗	✗	✗
Cholesterol Screening		✗ (24 mo–4 yrs)	✗	✗
STD Screening				✗
Pelvic Exam				✗
Anticipatory Guidance	✓	✓	✓	✓
Injury Prevention	✓	✓	✓	✓
Violence Prevention	✓	✓	✓	✓
Sleep Positioning Counseling	✓ (up to 6 mo.)			
Nutrition Counseling	✓	✓	✓	✓
Dental				
Prophylaxis as necessary (Cleaning)	✓	✓	✓	✓
Provide anticipatory guidance for parent/guardian	✓	✓ (12 – 24 mo)	✓ (2-6 yrs)	
Assessment of fluoride status & appropriate counseling/ prescription	✓	✓ (12 – 24 mo)	✓ (2-6 yrs)	
Recommend reevaluation every six months	✓	✓ (12 – 24 mo)	✓ (2-6 yrs)	
Restorative treatment or referral to specialist as indicated by patient's needs	✓	✓ (12 – 24 mo)	✓ (2-6 yrs)	
Review fluoride status		✓ (12 – 24 mo)	✓ (2-6 yrs)	
Radiographic assessment as indicated by patient's needs		✓ (12 – 24 mo)	✓ (2-6 yrs)	
Provide pit and fissure sealants for primary & permanent teeth as indicated by individual patient's needs		✓ (12 – 24 mo)	✓ (2-6 yrs)	
Provide assessment of developing malocclusions (deformities)		✓ (12 – 24 mo)	✓ (2-6 yrs)	
Provide diagnosis and treatment for any oral diseases, habits or injuries		✓ (12 – 24 mo)	✓ (2-6 yrs)	

Key:
✓ = to be performed   ✗ = to be performed for patients at risk
s = subjective, by history   o = objective, by a standard testing method
s/o = objective at 12, 15, and 18 years old, subjective, by history for all other years