

Wyoming Immunization Requirements and Recommendations For Children Attending a Child Caring Facility or Preschool

Vaccines in gray are recommended. All others are required.

| Child's Age | Vaccine | Number of Doses | Footnote |
|-------------------------------|--------------------------------|-------------------------------|---|
| 6 mos. | DTaP | Dose 3 of 5 | 3 doses at 2, 4, and 6 months. |
| | Hib | Dose 3 of 4 or Dose 2 of 3 | If ActHIB, MenHibrix, Hiberix, or Pentacel vaccines used, it is 3 doses at 2, 4, and 6 mos. If PedvaxHIB is used, it is 2 doses at 2 and 4 mos.; a dose at age 6 mos. is not needed. |
| | Pneumococcal conjugate (PCV13) | Dose 3 of 4 | 3 doses at 2, 4, and 6 mos. |
| | Rotavirus* | Dose 2 of 2 or Dose 3 of 3 | If Rotarix is used, a 2–dose series at 2 and 4 mos. If RotaTeq is used, a 3–dose series at 2, 4, and 6 mos. *If any dose in the series was RotaTeq or vaccine product is unknown for any dose in the series, a total of 3 doses of RV vaccine should be administered. |
| 6 to 18 mos. | Hepatitis B | Dose 3 of 3 | 1st dose at birth, 2nd dose between 1-2 mos., 3rd dose between 6-18 mos. |
| | Polio (IPV) | Dose 3 of 4 | 1st dose at 2 mos., 2nd dose at 4 mos., 3rd dose at 6-18 mos.. |
| 6 mos.+ (annually) | Influenza | Dose 1 of 2 | If receiving for the first time, 2 dose should be administered at least 4 weeks apart. After initial dose, vaccine is needed annually. |
| | Influenza | Dose 2 of 2 | |
| 12 to 15 mos. | Hib | Dose 3 of 3 or Dose 4 of 4 | One booster dose (dose 3 or 4, depending on vaccine used in primary series). |
| | Pneumococcal conjugate (PCV13) | Dose 4 of 4 | |
| | MMR | Dose 1 of 2 | |
| | Varicella | Dose 1 of 2 | If a child has had chickenpox the parent, guardian, school nurse, or physician must provide a written statement for the student's permanent school record. |
| 12 - 23 mos. | Hepatitis A | Dose 1 of 2 | |
| 15 to 18 mos. | DTaP | Dose 4 of 5 | 4th dose between 15-18 mos., 5th dose at 4-6 yrs. *If a 4th (4th) dose of DTaP vaccine was administered on or after a child's 4th (4th) birthday, and at least six (6) mos. has passed since the 3rd (3rd) dose, a fifth (5th) dose is not required. |
| 18 mos.+ | Hepatitis A | Dose 2 of 2 | 2nd dose 6-18 mos. after the first dose |
| 4 to 6 yrs. | DTaP | Dose 5 of 5 | |
| | Polio (IPV) | Dose 4 of 4 | The 4th dose is not needed if the 3rd dose is given on or after the fourth birthday. |
| | MMR | Dose 2 of 2 | |
| | Varicella | Dose 2 of 2 | |