
CANCER IN WYOMNG

Alexia Harrist MD, PhD

State Epidemiologist and State Health Officer

Wyoming Department of Health



Leading Causes of Death in WY

Cause of Death	2015	2016
Heart Disease	1,020	1,038
Cancer	924	955
Accidents/Injuries	393	362
Chronic Respiratory Disease	365	327
Stroke	195	206
Alzheimer's	151	136
Diabetes	134	112
Chronic Disease	59%	60%
Cancer	19%	20%





WY 2015 Cancer Incidence:

Prostate = 352

Breast = 346 (340 Female, 6 Male)

Lung = 276 (145 Female, 131 Male)

Colorectal = 196

Melanoma = 148

TOTAL = 2,577




WY 2015 Cancer Mortality:


- •Lung = 213 (107 Female, 106 Male)
- •Colorectal = 90
- •III-Defined = 72
- •Breast = 71
- Pancreas = 67

TOTAL = 924


INCIDENCE COUNT - ALL CANCER SITES - 5 YEAR PERIODS




WY POPULATION




INCIDENCE COUNT - ALL CANCER SITES - 5 YEAR PERIODS



WY CANCER RATES - ALL SITES Per 100,000 Population




Incidence Rate per 100,000 Population of New Cancers in the United States, 2010-2014



Wyoming Rate = 410.7 (403.5-418.0) U.S. Rate = 452


Cancer Mortality Rate per 100,000 Population in the United States, 2010-2014




Wyoming Rate = 153.6 (149.2-158.1) U.S. Rate = 166

WY CANCER RATES - PROSTATE

Per 100,000 Population




Incidence Rate per 100,000 Population of Prostate Cancer in the United States, 2010-2014




Wyoming Rate = 108.5 (103.4-113.8) U.S. Rate = 115

WY CANCER RATES - BREAST (FEMALE)

Per 100,000 Population



WY BREAST CANCER DIAGNOSIS - EARLY v. LATE



Incidence Rate per 100,000 Population of Female Breast Cancer in the United States, 2010-2014




Female Breast, All Ages, All Races/Ethnicities, Female Rate per 100,000 women












Rate per 100,000 women

111.5 - 120.5 120.7 - 123.7 125.4 - 130.1 130.2 - 143.5 No data/data suppressed


Wyoming Rate = 113.5 (108.2-119.0) U.S. Rate = 124

WY CANCER RATES - LUNG

Per 100,000 Population




Incidence Rate per 100,000 Population of Lung Cancer in the United States, 2010-2014



Wyoming Rate = 46.2 (43.8-48.7) U.S. Rate = 61

WY CANCER RATES - COLORECTAL

Per 100,000 Population



Incidence Rate per 100,000 Population of Colorectal Cancer in the United States, 2010-2014

Rate of New Cancers in the United States


Colon and Rectum, All Ages, All Races/Ethnicities, Male and Female Rate per 100,000 people












Rate per 100,000 people


31.4 - 37.3 37.5 - 39.1 40.6 - 43.2 No data/data suppressed 43.4 - 50.0

Wyoming Rate = 35.8 (33.7-38.1)U.S. Rate = 40

WY CANCER RATES - PANCREAS




Incidence Rate per 100,000 Population of Pancreatic Cancer in the United States, 2010-2014




Wyoming Rate = 9.0 (7.9-10.1) U.S. Rate = 13

WY CANCER RATES - MELANOMA




WY CANCER INCIDENCE RATE - MELANOMA

Per 100,000 Population




Incidence Rate per 100,000 Population of Melanoma in the United States, 2010-2014




Wyoming Rate = 23.9 (22.2-25.8) U.S. Rate = 21

WY CANCER RATES – CERVICAL CANCER Per 100,000 Population




Percent of Adolescents Aged 13-17 Who Are Up-to-Date for HPV Vaccination — Wyoming, U.S., and Best State, 2016 (National Immunization Survey)



^{*} Indicates Wyoming is different from the U.S.

Incidence Rate per 100,000 Population of Cervical Cancer in the United States, 2010-2014



Wyoming Rate = 7.5 (6.1-9.1) U.S. Rate = 8

Relative Survival by Cancer Type: Wyoming, 2007-2015 (All Ages Combined)

Relative Survival	12 months	24 months	36 months	48 months	60 months
All Sites	81.20%	73.90%	69.50%	65.90%	62.70%
Bladder w/in situ	88.20%	80.00%	76.50%	71.50%	68.00%
Brain/CNS	56.30%	39.40%	34.80%	28.40%	23.40%
Breast (Female)	97.30%	94.70%	90.70%	87.70%	85.00%
Cervical (Malignant only)	86.00%	77.10%	70.80%	63.80%	60.90%
Colorectal	81.60%	70.80%	64.60%	58.90%	55.30%
Kidney	86.70%	80.20%	74.10%	69.90%	63.90%
Leukemia	74.40%	68.10%	60.80%	55.80%	51.90%
Lung	40.50%	24.40%	18.40%	15.00%	12.00%
Melanoma	98.30%	95.70%	94.20%	91.80%	90.70%
Non Hodgkin	82.10%	76.00%	74.40%	69.30%	64.50%
Oral Cavity	86.60%	76.60%	68.20%	63.20%	58.10%
Ovary	78.20%	64.50%	50.60%	41.90%	36.20%
Pancreas	29.30%	15.60%	10.00%	6.20%	6.20%
Prostate	99.30%	98.40%	97.40%	96.30%	94.70%
Thyroid	97.90%	97.60%	95.90%	95.90%	92.30%
Uterine	93.50%	88.00%	83.00%	80.10%	74.60%

Relative Survival by Cancer Type: Wyoming, 2007-2015 (All Ages Combined)

Relative Survival	Wyoming 60 months 2007-2015	United States 60 months 2001-2013
All Sites	62.70%	66.0%
Bladder w/in situ	68.00%	75.6%
Brain/CNS	23.40%	32.3%
Breast (Female)	85.00%	88.9%
Cervical (Malignant only)	60.90%	67.8%
Colorectal	55.30%	63.8%
Kidney	63.90%	71.2%
Leukemia	51.90%	53.9%
Lung	12.00%	18.7%
Melanoma	90.70%	89.5%
Non Hodgkin	64.50%	67.7%
Oral Cavity	58.10%	60.1%
Ovary	36.20%	45.8%
Pancreas	6.20%	8.4%
Prostate	94.70%	97.8%
Thyroid	92.30%	96.8%
Uterine	74.60%	80.5%

COSTS - WY

	INPATIENT		OUT PATIENT		
	# Discharges	Cost	# Discharges	Cost	
	J		<u> </u>		
HEART DISEASE	11,776	\$456,217,985.19	12,136	\$74,980,663.36	
CANCER	2,186	\$84,880,548.68	1,577	\$8,641,160.30	
COPD	5,926	\$206,450,855.54	5,748	\$29,495,289.47	
DIABETES	7,523	\$275,661,367.48	11,690	\$56,129,365.93	
STROKE	1,056	\$39,964,915.54	639	\$6,032,500.69	
TOTAL	28,467	\$1,063,175,672.43	31,790	\$175,878,979.75	

Risk Factors

Several risk factors associated with Cancer

- Physical Activity (too little)
- Poor Nutrition
- Smoking
- Age













Obesity by Year, WY BRFSS










Chris Madden. www.crismussen.co.uk

Current Smoking by Year, WY BRFSS





Percent of Women Aged 50-74 Who Had a Mammogram in the Past Two Years — Wyoming, U.S., and Best State, 2014-2016 (BRFSS)




^{*} Indicates Wyoming is different from the U.S.

Percent of Women Aged 50-74 Who Had a Mammogram in the Past Two Years by Household Income — Wyoming, 2016 (BRFSS)




Percent of Adults Aged 50-75 Who Fully Met USPSTF Recommendations for Colorectal Cancer Screening — Wyoming, U.S., and Best State, 2014-2016 (BRFSS)




^{*} Indicates Wyoming is different from the U.S.

Percent of Adults Aged 50-75 Who Fully Met USPSTF Recommendations for Colorectal Cancer Screening by Household Income — Wyoming, 2016 (BRFSS)





Percent of Women Aged 21-65 Who Had a Pap Test in the Past Three Years — Wyoming, U.S., and Best State (BRFSS)




^{*} Indicates Wyoming is different from the U.S.

Women Aged 21-65 Who Had a Pap Test in the Past Three Years by Household Income — Wyoming, 2016 (BRFSS)







Lung Cancer Screening – United States, 2010-2015

	2010 NHIS	2015 NHIS
Number Eligible for Screening	8,456,800	6,819,500
Number Receiving Screening	276,700	262,700
Percent Screened (Confidence Interval)	3.3 (2.3-4.7)	3.9 (2.4-6.2)


According to data from the National Health Interview Survey, only a small percentage of eligible smokers are receiving low-dose computed tomography for lung cancer screening






"What fits your busy schedule better, exercising one hour a day or being dead 24 hours a day?"

Prostate Cancer Mortality Rate per 100,000 Population in the United States, 2010-2014




Wyoming Rate = 1.8 (16.2-21.1) U.S. Rate = 20

Female Breast Cancer Mortality Rate per 100,000 Population in the United States, 2010-2014



Wyoming Rate = 19.0 (16.9-21.2) U.S. Rate = 21

Lung Cancer Mortality Rate per 100,000 Population in the United States, 2010-2014



Wyoming Rate = 36.7 (34.5-38.9) U.S. Rate = 45