

Wyoming Department of Health Communicable Disease Snapshot of Screening Recommendations

<u>At Risk Populations</u>	<u>HIV</u>	<u>Hepatitis B*</u>	<u>Hepatitis C</u>	<u>Syphilis</u>	<u>Chlamydia</u>	<u>Gonorrhea</u>	<u>TB</u>
1945-1965 birth cohort (Baby Boomers)	X		X				
As indicated by employment (e.g. healthcare or congregate settings)							X
Active tuberculosis (TB) disease	X						
Recipient of blood transfusions, blood components, or organ transplant prior to July 1992	X	X	X				
Born in Asia, Africa or South America	X	X					X
Born in Latin America, Caribbean, Africa, Eastern Europe, and Russia	X						X
Current or past long-term hemodialysis	X	X	X				
Current or past history of homelessness	X						X
Current or past history of injection drug use or other intranasal illicit drug use	X	X	X				X
Current or past resident of detention or correction facility	X		X		X	X	X
Hepatitis B or C positive sexual contact (as indicated)	X	X	X		X	X	
Household contact to Hepatitis B positive individuals	X	X					
History of prior sexually transmitted diseases (STD)	X				X	X	
HIV infected		X	X	X	X	X	X
Household contact of patients with active TB disease							X
Men who have sex with men (MSM)	X	X	X	X	X	X	
More than one sex partner in last 60 days	X			X	X	X	
New sex partner in the last 60 days	X				X	X	
Parents born in Asia, Africa, or South America	X	X					
Pelvic inflammatory disease (PID)	X				X	X	
Pregnant women	X	X		X	X	X	
Recipient of clotting factor or blood concentrate prior to 1987	X		X				
Sexual contact with a STD positive individual (as indicated)	X			X	X	X	
Sexually active males and females < 25 years old	X				X	X	
Symptoms consistent with disease (as indicated)	X	X	X	X	X	X	X
Consistently abnormal liver tests		X	X				
Travelers with prolonged stays in countries where TB is endemic							X
Underlying immunocompromising conditions/immunosuppressive (including chemo) therapy	X	X					X

DO NOT screen solely on risks listed; complete the respective risk assessment to determine appropriate screenings.

***Prior to Hepatitis B testing, check WYIR for vaccination status. Fully vaccinated patients do not need screened for Hepatitis B unless vaccine failure is suspected.**