

**For HIV, STDs, and Hepatitis test after each partner or exposure, or every 1-3 months for frequent partners or exposures.**

**Wyoming Department of Health Communicable Disease Snapshot of Screening Recommendations**

<u>At Risk Populations</u>	<u>HIV</u>	<u>Hep B*</u>	<u>Hep C</u>	<u>Syphilis</u>	<u>Chlamydia</u>	<u>Gonorrhea</u>	<u>TB</u>
<b>Sexually active individuals</b>	X				X	X	
<b>**Genital chlamydia/gonorrhea: history of &amp; never tested or history of genital sex since last tested</b>	X				X	X	
<b>Pharyngeal chlamydia/gonorrhea: history of &amp; never tested or history of oral sex since last tested</b>	X				X	X	
<b>Rectal chlamydia/gonorrhea: history of &amp; never tested or history of anal sex since last tested</b>	X				X	X	
<b>New sex partner in the last 60 days</b>	X				X	X	
<b>More than one sex partner in last 60 days</b>	X			X	X	X	
<b>Sexual contact with a STD positive individual (as indicated)</b>	X			X	X	X	
<b>*Men who have sex with men (MSM)</b>	X	X	X	X	X	X	
<b>History of prior sexually transmitted diseases (STD)</b>	X				X	X	
<b>Pelvic inflammatory disease (PID)</b>	X				X	X	
<b>*Pregnant women</b>	X	X	X	X	X	X	
<b>Sexual assault</b>	X	X	X	X	X	X	
<b>*Current or past history of injection or intranasal drug use</b>	X	X	X				X
<b>Current or past history of homelessness</b>	X	X	X				X
<b>Current or past resident of detention or correction facility</b>	X	X	X		X	X	X
<b>*HIV infected</b>		X	X	X	X	X	X
<b>1945-1965 birth cohort (Baby Boomers)</b>	X		X				
<b>*Hepatitis B or C positive sexual contact (as indicated)</b>	X	X	X		X	X	
<b>*Household contact to Hepatitis B positive individuals</b>	X	X					
<b>*Current or past long-term hemodialysis</b>	X	X	X				
<b>*Recipient of blood transfusions, blood components, or organ transplant prior to July 1992</b>	X	X	X				
<b>*Recipient of clotting factor or blood concentrate prior to 1987</b>	X	X	X				
<b>*Consistently abnormal liver tests</b>		X	X				
<b>*Underlying immunocompromising conditions/immunosuppressive (including chemo) therapy</b>	X	X					X
<b>Born in Latin America, Caribbean, Africa, Eastern Europe, and Russia</b>	X						X
<b>*Born in Asia, Africa or South America</b>	X	X					X
<b>*Parents born in Asia, Africa, or South America</b>	X	X					
<b>Active tuberculosis (TB) disease</b>	X						
<b>Household contact of patients with active TB disease</b>							X
<b>Travelers with prolonged stays in countries where TB is endemic</b>							X
<b>As indicated by employment (e.g. healthcare or congregate settings)</b>							X
<b>*Symptoms consistent with disease (as indicated)</b>	X	X	X	X	X	X	X

\*Prior to Hepatitis B testing check WylR for vaccination status. Do not test fully vaccinated patients for Hepatitis B unless vaccine failure is suspected.

\*\*For genital chlamydia/gonorrhea testing a vaginal self-swab is preferred for individuals with a vagina and urine is preferred for those without a vagina.