

WYOMING MEDICAID

Preferred Drug List (PDL) February 7, 2024

Drug classes not included on this list are not managed through a Preferred Drug List (PDL).
 HOWEVER, THIS EXCLUSION IS NOT A GUARANTEE OF PAYMENT OR COVERAGE. Dosage limits and other requirements may apply.
 Drugs new to market are non-preferred until a clinical review has been completed. PA criteria will apply to both the pediatric population,
 as well as the adult population for those plans where PA/PDL limits are allowed
 Unless otherwise noted on the PDL, generic substitution is mandatory.

Yellow highlighted items below indicate new changes to the PDL. Red font indicates quantity/dosage limits apply. *Indicates BRAND is Preferred. May Use DAW 5.
 Contact the Change Healthcare PA Helpdesk @ 877-207-1126 for prior authorization if client has primary insurance that will not cover the brand name medication.

Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at <http://wymedicaid.org> for additional criteria.

| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE! PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> | | |
|---------------------------------------|--|---|---|--|--|----------------------|
| ADDICTION | BUPRENORPHINE COMBINATIONS | | Client must have a diagnosis of opioid dependence or abuse. This is not to be used for the treatment of chronic pain. Prior authorization will be required before any narcotic, benzodiazepine, or carisoprodol prescription will be allowed between fills. Prior authorization will be required before any short-acting stimulant prescription from any doctor other than the prescriber of buprenorphine or Suboxone, will be allowed between fills. Oral buprenorphine will be approved for clients with a documented allergy to naloxone. Please submit PA requests on the "Oral Buprenorphine/Naloxone or Oral Buprenorphine" PA form available at www.wymedicaid.org . Dosage limits apply Prior authorization will be required for doses >24mg | buprenorphine (oral) buprenorphine/naloxone film (BRAND IS PREFERRED) ZUBSOLV | | |
| | | buprenorphine/naloxone tablets SUBOXONE FILM* | | | | |
| | NALOXONE | | | | Kloxxado, naloxone products, and Narcan nasal spray will be limited to one fill per 180 days without prior authorization. Naloxone formulations available in quantities of 10ml will require prior authorization. | naloxone nasal spray |
| | KLOXXADO naloxone NARCAN NASAL SPRAY | | | | | |
| | NALTREXONE | | Client must have a diagnosis of alcohol or opioid dependence. Prior authorization will be required before any narcotic, carisoprodol, or benzodiazepine prescription will be allowed between fills. Prior authorization will be required before a short-acting stimulant prescription from any doctor other than the prescriber of naltrexone or Vivitrol will be allowed between fills. | | | |
| | | naltrexone VIVITROL | | | | |
| ALLERGY / ASTHMA / COPD | ANTIHISTAMINES, MINIMALLY SEDATING | | Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | desloratadine CLARINEX RDT/SYRUP levocetirizine | | |
| | cetirizine fexofenadine loratadine | | | | | |
| | ANTIHISTAMINE/DECONGESTANT COMBINATIONS | | Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | CLARINEX-D | | |
| | cetirizine/pseudoephedrine fexofenadine/pseudoephedrine loratadine/pseudoephedrine | | | | | |
| | ANTICHOLINERGIC BRONCHODILATORS | | Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Spiriva 5 day STARTER package will be allowed one (1) time per recipient | TIOTRIPIUM BROM (use brand) TUDORZA YUPELRI | | |
| | ATROVENT HFA INCRUSE ELLIPTA Ipratropium SPIRIVA HANDIHALER SPIRIVA RESPIMAT | | | | | |
| | ANTICHOLINERGIC COMBINATION AGENTS | | Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. **Will also require the diagnosis of COPD. | BEVESPI BREZTRI DUAKLIR TRELEGY | | |
| | ANORO ELLIPTA** COMBIVENT STIOLTO | | | | | |
| | LEUKOTRIENE MODIFIERS | | Trial and failure of preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. | zafirlukast | | |
| montelukast | | | | | | |
| LONG ACTING BRONCHODILATORS | | Trial and failure of two (2) preferred agents greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. | BROVANA | | | |
| arformoterol SEREVENT STRIVERDI | | | | | | |
| NASAL ANTIHISTAMINES | | Trial and failure of preferred agent greater than or equal to 90 days in the last 12 months will be required before approval can be given for a non-preferred agent. | azelastine 0.15% DYMISTA (use separate agents) olopatadine 0.6% RYALTRIS | | | |
| azelastine 0.1% | | | | | | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | | |
|--|--|---|--|--|------------------|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> | |
| ALLERGY / ASTHMA / COPD continued | NASAL STEROIDS | | Trial and failure of two (2) preferred agents greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Budesonide will be approved for pregnancy. | DYMISTA (use separate agents) OMNARIS QNASL XHANCE ZETONNA | |
| | budesonide | | | | |
| | flunisolide fluticasone mometasone | | | | |
| | SHORT ACTING BRONCHODILATORS - INHALERS | | Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Prior authorization will be required after a total of 12 albuterol inhalers are dispensed within 365 days. Minimum day supply of 16 days is required | levalbuterol (BRAND IS PREFERRED) PROAIR DIGIHALER PROVENTIL HFA | |
| | albuterol HFA PROAIR RESPICLICK VENTOLIN HFA XOPENEX HFA* | | | | |
| | STERIOD INHALANTS | | Trial and failure of two (2) preferred agents greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. *Fluticasone HFA and Asmanex HFA will be approved for pediatric clients 8 years of age or younger. Alvesco will be approved for a history of oral thrush with steroid inhalants. | AIRDUO DIGIHALER ALVESCO ARMONAIR ASMANEX HFA* fluticasone HFA* QVAR REDIHALER | |
| | AIRDUO RESPICLICK | | | | |
| | ARNUITY ELIPTA ASMANEX TWISTHALER budesonide suspension PULMICORT FLEXHALER | | | | |
| | STERIOD COMBINATION AGENTS | | Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. **Will also require the diagnosis of COPD or uncontrolled asthma. Advair 7 and 14-day STARTER package will be allowed one (1) time per recipient. | fluticasone/vilanterol (use preferred agent) fluticasone/salmeterol 55-14/113-14/232-14 fluticasone/salmeterol 100-50/250-50/500-50 (BRAND IS PREFERRED) TRELEGY | |
| | BREO ELLIPTA** DULERA SYMBICORT* | | | | |
| WIXELA | | | | | |
| EPINEPHRINE | | | | AUVI-Q (use preferred agent) EPI-PEN (use preferred agent) | |
| epinephrine auto-injector pen | | | | | |
| EOSINOPHILIC ASTHMA AGENTS | | | *Approval for these agents will require additional clinical criteria which can be found on the Additional Therapeutic Criteria Chart | FASENRA* NUCALA* TEZSPIRE | |
| | | DUPIXENT XOLAIR | | | |
| ARTHRITIS | IMMUNOMODULATORS | | Client must have diagnosis of AS prior to approval of a preferred agent. To receive a non-preferred agent, client must have a diagnosis of AS and a 56-day trial and failure of both preferred agents. **Cimzia will be allowed for clients that are pregnant or breast-feeding Quantity Limits apply for all diagnoses: Enbrel 25mg - limited to 10 per month Enbrel 50mg - limited to 5 per month Humira 20mg - limited to 10 per month Humira 40mg - limited to 5 per month | CIMZIA** COSENTYX REMICADE RINVOQ SIMPONI TALTZ XELJANZ/XR | |
| | ANKYLOSING SPONDYLITIS (AS) | | | | |
| | | | ENBREL HUMIRA | | |
| | JUVENILE IDIOPATHIC ARTHRITIS (JIA) | | Client must have diagnosis of JIA prior to approval of a preferred agent. To receive a non-preferred agent, client must have a diagnosis of JIA and a 56-day trial and failure of both preferred agents. | ACTEMRA ILARIS ORENCIA XELJANZ/XR | |
| | | | | | ENBREL HUMIRA |
| | PSORIATIC ARTHRITIS (PA) | | Client must have diagnosis of PA prior to approval of a preferred agent. To receive a non-preferred agent, client must have a diagnosis of PA and a 56-day trial and failure of two of the three preferred agents. *Otezla starter pack is non-preferred **Cimzia will be allowed for clients that are pregnant or breast-feeding | CIMZIA** COSENTYX ORENCIA REMICADE RINVOQ SIMPONI STELARA TALTZ TREMIFYA XELJANZ/XR | |
| | | ENBREL HUMIRA OTEZLA* | | | |
| RHEUMATOID ARTHRITIS (RA) | | Client must have diagnosis of RA and a 56-day trial and failure of methotrexate prior to approval of a preferred agent. To receive a non-preferred agent, client must have a diagnosis of RA and a 56-day trial and failure of both preferred agents. *Cimzia will be allowed for clients that are pregnant or breast-feeding **See Dermatology criteria for Atopic Dermatitis approval | ACTEMRA CIMZIA* KEVZARA KINERET OLUMIANT ORENCIA REMICADE RINVOQ** RITUXAN SIMPONI XELJANZ/XR | | |
| | | | | ENBREL HUMIRA | |
| CONVULSIONS | INTERMITTENT, STEREOTYPIC SEIZURE EPISODES | | *Nayzilam will be allowed for patients 12 years of age and older | | |
| | diazepam gel NAVZILAM* VALTOCO | | | | |
| | ORAL ANTICONVULSANTS | | Preferred agents with clinical criteria will be limited to FDA approved indications related to seizures and epilepsy. Non-preferred agents require 30 day trial and failure of two preferred agents prior to approval. For indications not related to seizures and epilepsy, please refer to the Additional Therapeutic Criteria chart at www.wymedicaid.org . **Please refer to the Additional Therapeutic Criteria chart at www.wymedicaid.org for specific requirements. | APTIOM (use preferred agent) BRIVIACT (use preferred agent) clobazam** DIACOMIT** FINTEPLA** levetiracetam ER OXTELLAR (use preferred agent) TROKENDI XR (use preferred agent) XCOPRI VIMPAT (tablets) zonisamide oral susp. (use preferred agent) | |
| carbamazepine divalproex FELBAMATE fosphenytoin lacosamide (tablets) lamotrigine/XR levetiracetam oxcarbazepine phenytoin subvenite valproate/valproic acid VIMPAT (suspension) zonisamide | | BANZEL (tablets only) clonazepam EPIDIOLEX gabapentin pregabalin topiramate/ER sprinkle caps | | | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|--|---|---|---|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| CROHN'S | IMMUNOMODULATORS | | Client must have diagnosis of Crohn's prior to approval of the preferred agent. To receive a non-preferred agent, client must have a diagnosis of Crohn's and a 56-day trial and failure of the preferred agent. * Refer to Additional Therapeutics Clinical Criteria Chart for more information **Cimzia will be allowed for clients that are pregnant or breast-feeding | CIMZIA** ENTYVIO* REMICADE RINVOQ STELARA TYSABRI (additional criteria applies) |
| | | HUMIRA | | |
| DERMATOLOGY | BENZOYL PEROXIDE/CLINDAMYCIN COMBOs | | Clients must be 12 to 20 years of age. Requires prior authorization for clients less than 12 years of age. Acne combinations are limited to clients under the age of 21. | ACANYA (use preferred agent) ONEXTON (use preferred agent) |
| | | clindamycin/benzoyl peroxide 1-5% clindamycin/benzoyl peroxide 1.2-5% (Refrig) | | |
| | ISOTRETINOIN | | | ABSORICA (use preferred agents) |
| | AMNESTEEM CLARAVIS isotretinoin ZENATANE | | | |
| | CORTICOSTEROIDS - STEP 1 AGENTS C=CREAM; G=GEL; L=LOTION; O=OINTMENT | | Trial and failure of two preferred agents greater than or equal to 14 days in the last 90 days. | PANDEL TEXACORT 2.5% (S) |
| | LOW POTENCY | | | |
| | alclometasone desonide* fluocinolone 0.01% hydrocortisone butyrate 0.1% (C) hydrocortisone 1%, 2.5% (C,L,O) | | *Cream, ointment, and lotion formulations of Desonide are preferred. | |
| | MEDIUM POTENCY | | Trial and failure of two preferred agents greater than or equal to 14 days in the last 90 days. | Clocortolone Pivalate flurandrenol fluticasone 0.05% (L) hydrocortisone butyrate 0.1% (O) triamcinolone 0.05% (O) |
| | betamethasone valerate desoximetasone 0.05%, 0.25% (C) fluocinolone 0.025% fluticasone 0.05% (C) mometasone SYNALAR 0.025% (C, O) triamcinolone 0.025%, 0.1% | | | |
| | HIGH POTENCY | | Trial and failure of two preferred agents greater than or equal to 14 days in the last 90 days. | APEXICON 0.05% (C) amcinonide 0.1% (C,L,O) augmented betamethasone 0.05% (G,L,O) clobetasol 0.05% (L) desoximetasone 0.05%, 0.25% (G,O) diflorasone 0.05% (C) fluocinonide 0.1% (C) halcinonide 0.1% (C) HALOG 0.1% (O) |
| betamethasone dipropionate clobetasol/E 0.05% (C,G,O,S) diflorasone 0.05% (O) fluocinonide flurandrenolide fluticasone 0.005% (O) halobetasol TOPICORT 0.025% (C) triamcinolone 0.5% ULTRAVATE 0.05% (C,O) | | | | |
| IMMUNOMODULATORS - STEP 2 AGENTS | | To receive a step 2 agent : Trial and failure of a preferred medium or high potency topical corticosteroid greater than or equal to a 21 day trial in the last 90 days. Exceptions will be made for application to the face and for clients age 12 and under, a trial and failure of a preferred low potency corticosteroid greater than or equal to a 21 day trial in the last 90 days will be required. | pimecrolimus (brand preferred) | |
| | ELIDEL tacrolimus | | | |
| PHOSPHODIESTERASE 4 INHIBITOR - STEP 3 AGENT | | To receive a step 3 agent : Trial and failure of a preferred step 2 agent (immunomodulator) greater than or equal to a 21 day trial within the last 30 days. | EUCRISA | |
| ATOPIC DERMATITIS | | *Trial and failure of a preferred step 2 agent (immunomodulator) greater than or equal to a 21 day trial within the last 30 days required. Dupixent requires member be at least 6 months of age or older. No high-potency steroid trial will be necessary. **Trial and failure of all criteria to receive a step 3 agent as defined above including medium/high potency topical corticosteroid, preferred step 2 immunomodulator AND 56-day trial and failure of a preferred biologic for Atopic Dermatitis in Step 3 will be required for approval of the non-preferred agents. | ADBRY** CIBINQO** OPZELURA** RINVOQ** | |
| | DUPIXENT* | | | |
| PLAQUE PSORIASIS (PP) | | Client must have diagnosis of PP prior to approval of a preferred agent (Enbrel, Humira, or Otezla). To receive a non-preferred agent, client must have a diagnosis of PP and a 56-day trial and failure of two of the three preferred agents. *Sotyktu requires diagnosis of moderate to severe PP and 56 day trial and failure of Humira. **Cimzia will be allowed for clients that are pregnant or breast-feeding | CIMZIA** COSENTYX ILUMYA REMICADE SILIQ SKYRIZI STELARA TALTZ TREMIFYA | |
| | ENBREL HUMIRA OTEZLA SOTYKTU* | | | |
| SCABICIDES/PEDEICULICIDES | | Trial and failure of a preferred agent in the last 12 months. | malathion lotion NATROBA spinosad (BRAND IS PREFERRED) | |
| permethrin VANALICE | | | | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|--|--|---|---|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| DIABETES | DIABETES AGENTS | | | metformin SR 24HR osmotic release (use preferred agent) metformin SR 24HR modified release (use preferred agent) |
| | BIGUANIDES | | | |
| | metformin/ER | | | |
| | α–GLUCOSIDASE INHIBITORS | | Trial and failure of metformin and a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | miglitol |
| | acarbose | | | |
| | MEGLITINIDES | | Trial and failure of metformin and a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | repaglinide |
| | nateglinide | | | |
| | THIAZOLIDINEDIONES | | Trial and failure of metformin and a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | ACTOPLUS MET (use separate agents) |
| | pioglitazone | | | |
| | SULFONYLUREAS | | Trial and failure of metformin and a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | |
| | glimepiride/ER glipizide/ER glyburide/ER | | | |
| | DIPEPTIDYL PEPTIDASE 4 (DPP-4) INHIBITORS | | Trial and failure of metformin greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a preferred agent. A 90 day trial of failure of the preferred agent is required before approval can be given for a non-preferred agent. | alogliptin GLYXAMBI (use separate preferred agents) QTERN (use separate preferred agents) STEGLUJAN (use separate preferred agents) |
| | JANUVIA ONGLYZA TRADJENTA | | | |
| | DPP-4 INHIBITOR COMBO AGENTS | | Trial and failure of metformin greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a preferred agent. A 90 day trial of failure of the preferred agent is required before approval can be given for a non-preferred agent. | alogliptin/metformin alogliptin/pioglitazone (use separate preferred agents) JENTADUETO XR saxagliptin/metformin (use brand) |
| | JANUMET/XR JENTADUETO KOMBIGLYZE/XR | | | |
| | INCRETIN MIMETICS (GLP-1 RECEPTOR AGONISTS) | | Trial and failure of metformin greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a preferred agent unless ASCVD or risk factors are present, in which case the trial of metformin is waived. A 90 day trial of failure of the preferred agent is required before approval can be given for a non-preferred agent. *Rybelsus requires documentation of inability to use injectable agents. Dosage Limits Apply: Ozempic: 2mg/week Victoza: 1.8mg/day | BYDUREON MOUNJARO OZEMPIC* SOLIQUA RYBELSUS* (additional criteria applies) XULTOPHY (use separate preferred agents) |
| | BYETTA TRULICITY VICTOZA | | | |
| | SGLT2 INHIBITORS | | Trial and failure of metformin greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a preferred agent unless there is a diagnosis of ASCVD, CKD, or heart failure, in which case the trial of metformin will be waived. A 90 day trial and failure of a preferred agent is required before approval can be given for a non-preferred agent. | GLYXAMBI (use separate preferred agents) QTERN (use separate preferred agents) SEGLUROMET (use separate preferred agents) STEGLATRO STEGLUJAN (use separate preferred agents) SYNJARDY XR (use separate preferred agents) TRIJARDY XR (use separate preferred agents) |
| | FARXIGA INVOKAMET INVOKANA JARDIANCE SYNJARDY XIGDUO XR | | | |
| | FAST-ACTING INSULIN | | Prior authorization will be required when using two different delivery forms of the same type of insulin concurrently. | ADMELOG (use preferred agent) FIASP (use preferred agent) insulin lispro (use preferred agents) LYUMJEV |
| HUMALOG HUMALOG 75/25 HUMALOG JR. HUMALOG MIX NOVOLOG MIX | | | | |
| LONG-ACTING INSULIN | | Prior authorization will be required when using two different delivery forms of the same type of insulin concurrently. | BASAGLAR (use preferred agent) Insulin Glargine (use preferred agent) Insulin Degludec SOLIQUA TOUJEO (use preferred agent) TRESIBA* (use preferred agent) XULTOPHY (use separate preferred agents) | |
| LANTUS SOLOSTAR* LANTUS vial LEVEMIR | | | | |
| DIABETIC METERS/TEST STRIPS | | Quantity limits apply: Insulin Dependent Clients: 10 strips/day Non-Insulin Dependent Clients: 4 strips/day Clients are limited to 1 meter/365 days | ALL OTHER METERS AND TEST STRIPS | |
| FREESTYLE (strips only) FREESTYLE FREEDOM FREESTYLE FREEDOM LITE FREESTYLE INSULINX FREESTYLE PRECISION NEO B FREESTYLE SIDEKICK II ONE TOUCH ULTRA II ONE TOUCH ULTRA MINI ONE TOUCH ULTRA BLUE ONE TOUCH VERIO ONE TOUCH VERIO FLEX ONE TOUCH VERIO REFLECT ONE TOUCH VERIO IQ PRECISION XTRA | | | | |
| EXTERNAL DIABETIC DEVICES | | | | |
| OMNIPOD DASH OMNIPOD CLASSIC OMNIPOD 5 | | | | |
| CONTINUOUS BLOOD GLUCOSE MONITORS | | Prior authorization will be required to verify if the client is injecting insulin daily. Monitors will also be limited to the labeled age. | GUARDIAN MINIMED | |
| DEXCOM G6 DEXCOM G7 FREESTYLE LIBRE 2 FREESTYLE LIBRE 3 | | | | |
| ACUTE HYPOGLYCEMIA AGENTS | | | GVOKE (use preferred agent) | |
| BAOSIMI ZEGLALOGUE (autoinjector) | | | | |
| FIBROMYALGIA | FIBROMYALGIA | | | |
| amitriptyline cyclobenzaprine duloxetine | | gabapentin | pregabalin SAVELLA tablets (savella titration pak will not be covered) | |
| | | | Clients will not be allowed to take gabapentin and pregabalin concurrently | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | | |
|--|--|--|---|--|--|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> | |
| GASTROINTESTINAL | BOWEL EVACUANTS | | | CLENPIQ (use preferred agents) GAVILYTE H (use preferred agents) POLY-PREP (use preferred agents) SUFLAVE SUTAB | |
| | GAVILYTE G, N GOLYTELY MOVIPREP PEG 3350 SOLUTION SUPREP | | | | |
| | CHRONIC IDIOPATHIC CONSTIPATION | | AMITIZA LINZESS TRULANCE | Client must have a diagnosis of chronic idiopathic constipation to receive a preferred agent. To receive a non-preferred agent, the client must have a diagnosis of chronic idiopathic constipation and a 30-day trial and failure of a preferred agent within the last 12 months. | MOTTEGRITY |
| | DIGESTIVE ENZYMES | | | Prior authorization required. | PERTZYE VIOKACE |
| | CREON ZENPEP | | | | |
| | IRRITABLE BOWEL SYNDROME WITH CONSTIPATION | | AMITIZA LINZESS TRULANCE | Client must have a diagnosis of Irritable Bowel Syndrome (IBS) with constipation. | |
| | MESALAMINE | | | Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | mesalamine DR tab 800mg (BRAND IS PREFERRED) mesalamine DR tab 1.2gm (BRAND IS PREFERRED) mesalamine ER cap 0.375gm (BRAND IS PREFERRED) mesalamine sup 1000mg SFRWOWASA |
| | APRISO* LIALDA* mesalamine 400mg DR capsule mesalamine enema PENTASA | | | | |
| | OPIOID-INDUCED CONSTIPATION AGENTS | | AMITIZA | Client must have a diagnosis of opioid-induced constipation and a three (3) month trial and failure of a stool softener to receive the preferred agent. To receive the non-preferred agent, the client must have a diagnosis of opioid-induced constipation, a three (3) month trial and failure of a secretory agent, and a three (3) month trial and failure of the preferred agent. | MOVANTIK* RELISTOR SYMPROIC |
| | | | | *Movantik will be approved for a diagnosis of cancer or for clients in hospice or palliative care. | |
| PREGNANCY INDUCED NAUSEA/VOMITING | | | | | |
| BONJESTA DICLEGIS | | | | | |
| PROTON PUMP INHIBITORS | | | Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. PREVACID solutabs will be approved for children less than or equal to 8 years of age. | amox/clarith/lanso pack (use separate agents) DEXILANT dexlansoprazole esomeprazole omeprazole 20.6mg capsules (use preferred) omeprazole tablets (use preferred agent) omeprazole/sodium bicarbonate OMECLAMOX (use separate agents) PREVACID solutabs* rabeprazole TALICIA (use separate agents) VIMOVO (use separate agents) | |
| lansoprazole capsules omeprazole capsules pantoprazole | | | | | |
| GOUT | COLCHICINE | | | MITIGARE (use preferred agent) | |
| | colchicine (tablets) | | | | |
| XANTHINE OXIDASE AND URAT1 INHIBITORS | | | Trial and failure of the preferred agent greater than or equal to a 60 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | ULORIC* | |
| allopurinol | | | | | |
| HEMATOLOGY | LOW MOLECULAR WEIGHT HEPARIN (LMWH) | | Prior authorization will be required for the 300mg/3ml strength. | FRAGMIN (use preferred agent) enoxaparin 300MG/3ML | |
| | enoxaparin | | | | |
| | DIRECT THROMBIN INHIBITOR | | PRADAXA | Client must have diagnosis of non-valvular atrial fibrillation and relative contraindication to warfarin for approval, treatment for deep vein thrombosis (DVT) or pulmonary embolism (PE), or for the reduction in the risk of recurrence of DVT and PE after initial therapy. | |
| | | | | | |
| SELECTIVE FACTOR XA INHIBITOR | | XARELTO 2.5mg* (use preferred) | *To receive Xarelto 2.5mg, client must have a diagnosis of chronic coronary artery disease or peripheral artery disease with the need to reduce risk of major cardiovascular events | ELIQUIS (starter pack) SAVAYSA (use preferred agent) | |
| ELIQUIS XARELTO 10mg, 15mg, 20mg, and starter pack | | | | | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|--|---|---|--|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| HEMATOLOGY continued | CPTP DERIVATIVES | | Client must have a diagnosis of acute coronary syndrome, history of myocardial infarction, or history of stroke and transient ischemic attack. | |
| | | BRILINTA | | |
| | PAR-1 ANTAGONIST | | Client must have diagnosis of reduction of thrombotic cardiovascular events with a history of myocardial infarction (MI) or with peripheral arterial disease (PAD). Must be used in conjunction with aspirin or clopidogrel. | |
| | | ZONTIVITY | | |
| | ANTIHEMOPHILIC FACTOR VIII | | | ALTUVIIO KOVALTRY |
| | | ADVATE ADYNOVATE AFSTYLA ELOCTATE ESPEROCT HEMOFIL M HEMLIBRA JIVI KOATE/KOATE-DVI KOGENATE FS/BIO-SET NOVOEIGHT NUWIQ OBIZUR RECOMBINATE XYNTHA/SOLOFUSE | | |
| | COAGULATION FACTOR IX | | | |
| | ALPHANINE SD ALPROLIX BENEFIX IDELVION IXINITY REBINYN RIXUBIS | | | |
| ANTIHEMOPHILIC FACTOR/VWF | | | | |
| | ALPHANATE HUMATE-P VONVENDI WILATE | | | |
| ERYTHROPOIESIS STIMULATING AGENTS | | | | ARANESP PROCRIT |
| | EPOGEN MIRCERA RETACRIT | | | |
| SICKLE CELL ANEMIA | | | | |
| | DROXIA SIKLOS | | | |
| HEPATITIS C | DIRECT ACTING ANTIVIRALS | | Limited to FDA approved indication. Prior authorization will be required prior to use of preferred agents. Please submit PA requests on the Hepatitis CPA form available at www.wymedicaid.org . | EPCLUSA (use preferred agent) HARVONI SOVALDI VOSEVI** ZEPATIER |
| | | sofosbuvir/velpatasvir MAVYRET | | |
| HIDRADENITIS SUPPURATIVA | IMMUNOMODULATORS | | Humira will not be covered as a first line agent for the diagnosis of hidradenitis suppurativa. | |
| | | HUMIRA | | |
| HORMONES | GnRH ANTAGONISTS | | *Please refer to the Additional Therapeutic Criteria chart at www.wymedicaid.org for specific requirements. | ORILISSA |
| | MYFEMBREE ORIAHNN | | | |
| | GROWTH HORMONE | | | HUMATROPE NGENLA OMNITROPE SAIZEN SEROSTIM SKYTROFA SOGROYA ZOMACTON |
| | | GENOTROPIN NORDITROPIN NUTROPIN AQ | | |
| | TESTOSTERONE TOPICAL GELS | | Testosterone agents are only allowed for diagnosis of hypogonadism or insufficient | ANDRODERM (use preferred agent) FORTESTA (use preferred agent) JATENZO (use preferred agent) TESTOPEL (use preferred agent) testosterone gel (use preferred agent) testosterone solution (use preferred agent) XYOSTED (use preferred agent) |
| | ANDROGEL* TESTIM GEL | | | |
| THYROID HORMONES | | Ermeza will be covered with confirmed diagnosis of dysphagia. | | THYQUIDITY TIROSINT |
| | ARMOUR THYROID LEVOXYL levothyroxine (tablets) LEVO-T liothyronine SYNTHROID UNITHROID | ERMEZA | | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|--|--|-------------------|---|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| HORMONES continued | ORAL CONTRACEPTIVES | | | |
| | afirmelle | | | alyacen 1-35, 7/7/7 |
| | altavera | | | aranelle |
| | amethia | | | BALCOLTRA |
| | amethyst | | | balziva |
| | apri | | | brillyn |
| | ashlyna | | | drospr/ethinyl estradiol/levomefolate |
| | aubra/EQ | | | enpresse |
| | aurovela 1-20/FE 1-20, 1-35 | | | ethynodiol/ethinyl estradiol |
| | aviane | | | FALESSA KIT |
| | ayuna | | | fayosim |
| | azurette | | | kaitlib FE chew |
| | blisovi 1-20 FE, 1.5-30 FE | | | layolis FE chew |
| | bekyree | | | levonest |
| | beyaz | | | levonorgest/ethinyl estradiol/LO (84-7) |
| | camrese/LO | | | levonorgest/ethinyl estradiol 0.15- |
| | chateal/EQ | | | MINASTRIN FE chew* |
| | CHARLOTTE 24 FE chew | | | noreth/ethinyl estradiol/FE chew 0.8/25 |
| | cyred | | | nortrel |
| | dasetta 1-35, 7/7/7 | | | phiith |
| | daysee | | | rivelsa |
| | deso/ethinyl estradiol | | | QUARTETTE |
| | drospr/ethinyl estradiol | | | SAFYRAL |
| | elinest | | | TAYTULLA |
| | enskyce | | | tilia FE |
| | estarylla | | | tri-legest FE |
| | falmina | | | TRIVORA |
| | finzala FE chew | | | TWIRLA |
| | gianvi | | | tydemy |
| | halley FE 1/20, 1/35 | | | vyfemla |
| | iclevia | | | wera |
| | introvoile | | | wymzya FE chew |
| | isibloom | | | |
| | laimiess | | | |
| | jolessa | | | |
| | juleber | | | |
| | junel 1-20/FE, 1.5-30/FE | | | |
| | kalliga | | | |
| | kariva | | | |
| | kelnor | | | |
| | kurvelo | | | |
| | larin 1-20/FE, 1.5-30/FE | | | |
| | leena | | | |
| | lessina | | | |
| | levora | | | |
| | lo loestrin | | | |
| | loestrin FE | | | |
| | loryna | | | |
| | LOSEASONIQUE* | | | |
| | low-ogestrel | | | |
| | luteru | | | |
| | marlissa | | | |
| | melodetta | | | |
| | mibelas FE chew | | | |
| | microgestin 1-20/FE, 1.5-30/FE | | | |
| | mili | | | |
| | mono-linyah | | | |
| | natazia | | | |
| | NECON 0.5/35, 1/35, 1/50, 7/7/7, 10/11 | | | |
| | nikki | | | |
| | noreth/ethinyl estradiol/FE chw 0.4/35, 1/20 | | | |
| | noreth/ethinyl estradiol 1-20/FE | | | |
| | norgest/ethinyl estradiol/LO | | | |
| | norethindrone | | | |
| | nylia | | | |
| | nymyo | | | |
| | ocella | | | |
| | pimtreu | | | |
| | portia | | | |
| | previfem | | | |
| | reclipsen | | | |
| | safyral | | | |
| | SEASONIQUE* | | | |
| | setlakin | | | |
| | simliya | | | |
| | simpesse | | | |
| | sprintec | | | |
| | sronyx | | | |
| | syeda | | | |
| | tri-estaryl/LO | | | |
| | tri-femynor | | | |
| | tri-linyah | | | |
| | tri-marzia LO | | | |
| | tri-mili/LO | | | |
| | tri-sprintec/LO | | | |
| | tri-nymyo | | | |
| | tri-vyibra | | | |
| | velvet | | | |
| | vestura | | | |
| | vienva | | | |
| | violele | | | |
| | volnea | | | |
| | vyibra | | | |
| | yasmin-28 | | | |
| | YAZ | | | |
| | zumandimine | | | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|---|--|---|---|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| HYPERLIPIDEMIA | BILE ACID SEQUESTRANT | | Trial and failure of ALL preferred agents greater than or equal to six (6) months in the last 12 months will be required before approval can be given for a non-preferred agent. | WELCHOL |
| | cholestyramine/light colestipol | | | |
| | STATINS, LOW POTENCY | | Trial and failure of a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. If client's current medication therapy is contraindicated with the preferred statin(s) due to a drug-drug interaction, a non-preferred agent may be obtained with a prior authorization. Prior authorization will be required for clients under the age of 10. | fluvastatin/ER |
| | lovastatin pravastatin | | | |
| | STATINS, HIGH POTENCY | | Trial and failure of a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. If client's current medication therapy is contraindicated with the preferred statin(s) due to a drug-drug interaction, a non-preferred agent may be obtained with a prior authorization. Prior authorization will be required for clients under the age of 10. | EZALLOR LIVALO ZYPITAMAG |
| | atorvastatin rosuvastatin simvastatin | | | |
| | STATIN COMBINATIONS | | Trial and failure of a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Prior authorization will be required for clients under the age of 10. | ezetimibe/simvastatin (BRAND IS PREFERRED) |
| amlodipine/atorvastatin VYTORIN* | | | | |
| PCSK9-RELATED AGENTS | | PRALUENT | Client must have a diagnosis of homozygous familial hypercholesterolemia; have a diagnosis of heterozygous familial hypercholesterolemia or atherosclerotic cardiovascular disease AND not at goal with a maximum dose statin; or be intolerant to statin therapy. Approval for a non-preferred agent requires trial and failure of a preferred agent. | LEQVIO REPATHA |
| TRIGLYCERIDE LOWERING AGENTS | | fenofibrate 40/48/54/67/134/145/160/200mg gemfibrozil | Trial and failure of a preferred agent greater than or equal to a 90 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | fenofibric fenofibrate (43/50/120/130/150mg) icosapent LIPOFEN omega-3-acid VASCEPA |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | | |
|--|---|--|---|--|---|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> | |
| HYPERTENSION/ CARDIOLOGY | ANGIOTENSIN RECEPTOR BLOCKERS (ARBs) EDARBI irbesartan losartan olmesartan telmisartan valsartan | | Non-preferred ARBs will require a history of ALL preferred ARBs before approval | candesartan eprosartan 600mg | |
| | ARBs AND DIURETICS EDARBYCLOR irbesartan HCTZ losartan HCT olmesartan HCTZ valsartan HCTZ | | Non-preferred ARB/diuretic combinations will require a history of ALL preferred | candesartan HCTZ telmisartan HCTZ | |
| | ALPHA-BLOCKERS clonidine clonidine TD patches | | | | |
| | COMBINATION PRODUCTS | ENTRESTO | Client must be greater than one year of age and have a diagnosis of Congestive Heart Failure (CHF) NYHA Class II-IV. Angiotensin converting enzyme (ACE) inhibitors and angiotensin receptor blockers (ARBs) will not be allowed in combination with Entresto. | VERQUVO | |
| INFECTIOUS DISEASE | QUINOLONES ciprofloxacin levofloxacin ofloxacin | | Please refer to the Additional Therapeutic Criteria Chart located at http://www.wymedicaid.org/additional-therapeutic-criteria for Baxdela criteria. | moxifloxacin (use preferred agents) | |
| | DOXYCYCLINE doxycycline | | | DORYX (use preferred agent) | |
| | MINOCYCLINE minocycline/ER | | | minocycline 65mg and 115mg ER (use preferred agent) SOLODYN (use preferred agent) | |
| | INHALED TOBRAMYCIN KITABIS | | *Tobi Podhaler requires a 28 day trial of a preferred agent, as well as 28 days off of that same preferred agent prior to approval. Minimum day supply of at 56 days is required | BETHKIS inhaled tobramycin TOBI PODHALER (use preferred agent) | |
| | ANTI-RETROVIRALS APRETUDE BIKTARVY CIMDUO DELSTRIGO DOVATO EVOTAZ GENVOYA ODEFSEY PIFELTRO PREZCOBIX ritonavir tablets SYMFI/LO TRIUMEQ TROGARZO | CABENUVA* DESCOVY* TRUVADA* | | *Please refer to the Additional Therapeutic Criteria chart at www.wymedicaid.org for specific requirements. **Rukobia approval requires documentation of multi-drug resistance defined as failure of two medications from different classes. | JULUCA NORVIR RUKOBIA** STRIBILD (use separate agents) SUNLENCA SYMTOZA (use separate preferred agents) |
| | NSAIDs celecoxib diclofenac tablets etodolac FLECTOR* flurbiprofen ibuprofen indomethacin ketoprofen ketorolac meclofenamate meloxicam nabumetone naproxen oxaprozin piroxicam sulindac | | | Trial and failure of two (2) preferred agents each greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Dosing and quantity limits apply for ketorolac (limit 5days/34 days; max dose 40mg/day for oral tablets). | CALDOLOR (use preferred agent) diclofenac 1.3% patch (BRAND IS PREFERRED) diclofenac 1.5% soln. diclofenac 3% gel fenoprofen mefenamic acid NEOPROFEN (use preferred agent) |
| INFLAMMATION | ORAL CORTICOSTEROIDS budesonide cortisone acetate dexamethasone/intensol hydrocortisone methylprednisolone prednisolone prednisone | | | CELESTONE (use preferred agent) | |
| | NON-BENZODIAZEPINES BELSOMRA eszopiclone zaleplon zolpidem zolpidem ER | | Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Prior Authorization will be required for clients under the age of 18 *Quviviq requires trial and failure of two preferred agents with different mechanisms of action **Rozerem is non-preferred without a history of substance abuse Prior authorization will be required when a client is taking more than one insomnia agent concurrently. Dosage limits apply: zaleplon: 30mg/day zolpidem: 15mg/day | EDLUAR (additional criteria applies) DAYVIGO QUVIVIQ ROZEREM* zolpidem sublingual (additional criteria applies) | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | | |
|---|---|--|--|---|---|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> | |
| MENTAL HEALTH | ALZHEIMER'S AGENTS | | Client must have a diagnosis of dementia. Trial and failure of two (2) preferred agents greater than or equal to six (6) weeks WITHIN THE LAST 2 YEARS will be required before approval can be given for a non-preferred agent. One of the trials of preferred agents must be in the same class (NaSS, NDRI, SSRI, or SNRI) as the requested non-preferred agent. Trazodone, buspirone, fluvoxamine, MAO inhibitors, TCA's, bupropion IR, and venlafaxine IR do not require prior authorization but will not count towards meeting preferred therapy requirements. Clients will not be allowed to be on more than one antidepressant, including fluvoxamine, bupropion IR, and venlafaxine IR, at one time with the exception of mirtazapine or bupropion with a SSRI or SNRI. ***Trintellix requires trial and failure of two preferred agents in any class Clients five (5) years of age and younger will require prior authorization before approval. Dosage limits apply: bupropion ER/SR/XL: 450mg/day citalopram < 60 years of age: 60mg/day citalopram > 60 years of age: 30mg/day escitalopram: 30mg/day fluoxetine < 18 years of age: 90mg/day fluoxetine > 18 years of age: 120mg/day mirtazapine: 67.5mg/day paroxetine IR/CR < 18 years of age: 75mg/day paroxetine IR > 18 years of age: 90mg/day paroxetine CR > 18 years of age: 112.5mg/day sertraline: 300mg/day venlafaxine ER: 337.5mg/day | donepezil/ODT galantamine/ER memantine tablets/solution | donepezil 23mg (use preferred agent) memantine ER NAMZARIC (use separate agents) rivastigmine capsules/patches |
| | ANTIDEPRESSANTS | | | | |
| | NORADRENERGIC/SPECIFIC SEROTONERGICS (NaSS) | | | | NaSS |
| | mirtazapine tablets | | | | mirtazapine rapid dissolve tablets (use preferred agent) |
| | NOREPINEPHRINE/DOPAMINE REUPTAKE INHIBITORS (NDRI) | | | | NDRI |
| | bupropion ER/SR/XL | | | | APLENZIN AUVELITY FORFIVO XL* |
| | SELECTIVE SEROTONIN REUPTAKE INHIBITORS (SSRI) | | | | SSRI |
| | citalopram escitalopram fluoxetine capsules paroxetine IR/CR sertraline | | | | citalopram capsules fluoxetine tablets VIIBRYD |
| | SEROTONIN/NORPINEPHRINE REUPTAKE INHIBITORS (SNRI) | | | | SNRI |
| | duloxetine venlafaxine ER capsules | | | | desvenlafaxine FETZIMA venlafaxine ER tablets (use preferred agent) |
| ATYPICAL ANTIPSYCHOTICS | | | OTHER | | |
| ABILIFY MAINTENA ABILIFY ASIMTUFI aripiprazole tab/solution/ODT ARISTADA FANAPT** paliperidone INVEGA HAFYERA/SUSTENNA/TRINZA lurasidone olanzapine PERSERIS quetiapine* quetiapine ER RISPERDAL CONSTA risperidone SAPHRIS** VRAYLAR ziprasidone | | | *Quetiapine doses less than 100mg will require prior authorization <u>without</u> a diagnosis of mood disorder or major depressive disorder. For titration doses, contact the Change Healthcare Pharmacy Help Desk for an override. **Clients nine (9) years of age and younger will require a prior authorization to receive approval of Latuda and Saphris. Clients eighteen (18) years of age and younger will require a prior authorization to receive approval of Fanapt. ***Rexulti approval for MDD treatment requires concurrent antidepressant therapy as well as a trial and failure of aripiprazole or other preferred atypical antipsychotic indicated for the adjunct treatment of MDD. Trial and failure of two (2) preferred agents greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent unless otherwise specified. Prior authorization will be required for any client five (5) years of age or younger, or for any client taking both an injectable and oral dosage form of the same medication concurrently. Dosage limits apply: aripiprazole <13 years of age: 15mg/day aripiprazole ≥13 years of age: 30mg/day ABILIFY MAINTENA: 400mg per 26 days ARISTADA 441/662/882mg: 1 injection per 28 days ARISTADA 1064mg: 1 injection per 56 days ARISTADA INITIO: 1 injection per 365 days FANAPT: 24mg/day INVEGA HAFYERA: 1 injection per 6 months INVEGA SUSTENNA: 1 injection per 28 days INVEGA TRINZ: 1 injection per 84 days LATUDA 10-17 years of age: 80mg/day LATUDA >17 years of age: 160mg/day olanzapine <13 years of age: 10mg/day olanzapine ≥13 years of age: 20mg/day paliperidone: 12mg/day PERSERIS: 1 injection per 28 days quetiapine <13 years of age: 400mg/day quetiapine 13-17 years of age: 600mg/day quetiapine >17 years of age: 800mg/day risperidone <10 years of age: 3mg/day risperidone 10-17 years of age: 6mg/day risperidone >17 years of age: 16mg/day RISPERDAL CONSTA: 2 injections per 28 days SAPHRIS: 20mg/day ziprasidone ≤17 years of age: 120mg/day ziprasidone >17 years of age: 200mg/day | TRINTELIX*** | |
| SPECIAL ATYPICAL ANTIPSYCHOTICS | | | | | |
| clozapine/ODT | | | Dosage limits apply: 900mg/day | VERSACLOZ Suspension (use preferred agent) | |

**WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024**

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|--|--|---|--|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| MENTAL HEALTH continued | AMPHETAMINES LONG ACTING AMPHETAMINES ADDERALL XR amphetamine salts combo XR dextroamphetamine CR caps VYVANSE CAPSULES** | | Clients over the age of 17 must have a diagnosis for ADD, ADHD, (see ADD/ADHD criteria below), narcolepsy, obstructive sleep apnea, shift work sleep disturbance, MS fatigue (see MS Fatigue criteria below), or refractory depression (see refractory depression criteria below). For clients over the age of 17, they must meet the DSM-5 criteria for diagnosis of ADHD. These criteria include: • Five or more symptoms of inattention, present for at least 6 months, inappropriate for developmental level. OR • Five or more symptoms of hyperactivity and impulsivity, present for at least 6 months, to an extent that is disruptive and inappropriate for developmental level. AND • Symptoms must be present in two or more settings (home, school or work); and • There must be clear evidence that the symptoms interfere or reduce the quality of social, school or work functioning; and • The symptoms must not be better explained by another mental disorder. | AMPHETAMINES ADZENYS XR ODT DYANAVEL XR EVEKEO/ODT MYDAYIS PROCENTRA VYVANSE CHEWABLES ZENZEDI 2.5 AND 7.5MG TABLETS |
| | IMMEDIATE RELEASE AMPHETAMINES amphetamine salts combo dextroamphetamine tablets | | | Diagnosis of MS fatigue will require a fatigue severity scale score of 5.0, a 60-day trial of amantadine and discontinuation of medications that may contribute to drowsiness and fatigue. Diagnosis of refractory depression will require a 6-week trial and failure of an antidepressant (monotherapy) and continued concomitant use of an antidepressant with the stimulant. Prior Authorization will be required for clients under the age of 4. **Vyvanse will be approved for the diagnosis of binge-eating disorder for clients 18 years of age and older. Authorizations will be approved for 12 weeks. After initiation a one-year prior authorization of Vyvanse for this diagnosis will require additional documentation prior to approval. Claims will require Prior Authorization if clients have a history of the following: glaucoma, cardiac arrhythmias, arteriosclerosis, untreated hypertension, untreated hyperthyroidism, substance abuse, or current MAO inhibitor use. Trial and failure of two (2) preferred agents (each from a different class: methylphenidate and amphetamine) greater than or equal to a 30 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Two or more long-acting agents will not be allowed concurrently. Dosage limits apply: amphetamine salts combo XR: 60mg/day amphetamine salts combo: 60mg/day amphetamine salts combo (narcolepsy): 90mg/day DAYTRANA: 45mg/9 hour patch/day dextroamphetamine: 90mg/day dextroamphetamine CR: 90mg/day dexmethylphenidate: 30mg/day FOCALIN XR < 13 years of age: 45mg/day FOCALIN XR > 13 years of age: 60mg/day JORNAY PM: 100mg/day methylin/methylphenidate/ER: 90mg/day VYVANSE: 105mg/day |
| | METHYLPHENIDATES LONG ACTING METHYLPHENIDATES CONCERTA* dexmethylphenidate ER methylphenidate ER tablets | | | |
| | IMMEDIATE RELEASE METHYLPHENIDATES dexmethylphenidate methylphenidate chewables methylphenidate solution methylphenidate tablets | | | |
| | SELECTIVE ALPHA-ADRENERGIC AGONIST clonidine | | To obtain the non-preferred agent , client must meet the following criteria: Client must have a diagnosis of ADD or ADHD Prior authorization will be required for clients under the age of 4. To receive clonidine ER, clients must have completed a 14 day trial of clonidine IR with benefit in the previous 12 months. | clonidine ER |

**WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024**

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|---|---|---|--|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| MENTAL HEALTH continued | SELECTIVE NOREPINEPHRINE REUPTAKE INHIBITOR | | Atomoxetine: Clients must have a diagnosis for ADD, ADHD, narcolepsy, obstructive sleep apnea, shift work sleep disturbance, or refractory depression (see refractory depression criteria below). Diagnosis of refractory depression will require a 6-week trial and failure of an antidepressant (monotherapy) and continued concomitant use of an antidepressant with the stimulant. Prior Authorization required for clients under the age of 4. Claims will require Prior Authorization if clients have a history of the following: glaucoma, cardiac arrhythmias, arteriosclerosis, untreated hypertension, untreated hyperthyroidism, substance abuse, or current MAO inhibitor use. Qelbree: Clients must have a 30-day trial and failure of a preferred non-stimulant. Approval will be limited to a max dose of 400mg in clients 6-17 years of age, and 600mg for clients 18 years of age or older. Dosage limits apply: atomoxetine: 100mg/day | QELBREE |
| | | atomoxetine | | |
| MIGRAINE | MIGRAINE PROPHYLAXIS | | Trial and failure of both an anticonvulsant and a beta blocker (Step 1 agents) greater than or equal to three (3) months will be required before approval can be given for the step 2 agents. Concurrent use of Botox will not be approved. Nurtec will be limited to 16 tabs/30 days. | NURTEC |
| | STEP 1 AGENTS | | | |
| | beta blockers | divalproex topiramate | | |
| | STEP 2 AGENTS | | *Starting dose will be limited to 70mg **Approval for non-preferred agents requires trial and failure of a preferred agent along with the trial and failures described with Step 1 Agents' criteria above. | QULIPTA** |
| | ACUTE MIGRAINE TREATMENT | | | |
| STEP 1 AGENTS | | Trial and failure of two preferred agents will be required for approval of a non-preferred agent. Rizatriptan will be limited to clients 6 years of age or older Quantity limits apply: natriptan 1mg: 25 tabs/34 days natriptan 2.5mg: 10 tabs/34 days RELPAX 20mg: 20 tabs/34 days RELPAX 40mg: 14 tabs/34 days rizatriptan 5mg: 27 doses/34 days rizatriptan 10mg: 14 doses/34 days sumatriptan vials: 2 vials/34 days sumatriptan nasal 20mg: 6 bottles/34 days; 5mg: 12 bottles/34 days sumatriptan 25mg: 41 tabs/34 days sumatriptan 50mg: 20 tabs/34 days sumatriptan 100mg: 10 tabs/34 days | almotriptan ELYXB Sumatriptan-Naproxen Sodium TOSYMRA (use preferred agent) TROKENDI XR ZEMBACE (use preferred agent) zolmitriptan | |
| STEP 2 AGENTS | | Trial and failure of two triptan agents is required for approval of a Step 2 Agent. Trial and failure of two preferred triptan agents AND Nurtec will be required for approval of a non-preferred agent. Quantity limits apply: NURTEC 75mg: limited to 8 tabs/30 days REYVOW: 200mg/day or 1 tab/day, 4 tab/30 days | REYVOW JUBRELVY | |
| MOVEMENT DISORDERS | VMAT 2 INHIBITORS | | Quantily limits apply: AUSTEDO: limited to 4 tabs/day INGREZZA: limited to 4 tabs/day *Please refer to the Additional Therapeutic Criteria chart at www.wymedicaid.org for specific requirements. | |
| | AUSTEDO/XR* INGREZZA* TETRABENAZINE | | | |
| MULTIPLE SCLEROSIS | MS AGENTS | | Kesimpta, Lemtrada, Ocrevus, and Tysabri will be approved for highly active disease, please refer to the ATCC for additional information. Trial and failure of two preferred agents for at least 56 days (each from a separate class) will be required before approval can be given for a non-preferred agent. For Mavenclad, in addition to the above criteria, approval will be granted on a case-by-case basis. | AUBAGIO BAFIERTAM BRIUMVI EXTAVIA glatiramer (BRAND IS PREFERRED) GLATOPIA (use preferred agent) MAVENCLAD MAYZENT PLEGRIDY PONVORY TECFIDERA ZEPOSIA |
| | AVONEX BETASERON COPAXONE 20MG/ML* dimethyl fumarate REBIF teriflunomide VUMERITY | GILENYA KESIMPTA LEMTRADA OCREVUS TYSABRI | | |
| NARCOLEPSY | STIMULANTS | | Modafinil and Nuvigil: Client must be > 16 years of age. Client must have a diagnosis of narcolepsy, obstructive sleep apnea, shift work sleep disorder, Multiple Sclerosis (MS) Fatigue, or ADD/ADHD with a concurrent diagnosis of substance abuse. Diagnosis of MS will require a fatigue severity scale score of 5.0, a 60 day trial of amantadine AND discontinuation of medications that may contribute to drowsiness or fatigue. Clients will not be allowed to take two or more agents in this class concurrently | SUNOSI WAKIX XYREM |
| | NON-STIMULANTS | | | |
| NEUROPATHIC PAIN | GABAPENTIN | | Clients will not be allowed to take gabapentin and pregabalin concurrently Prior authorizations for perioperative pain will be approved for gabapentin OR pregabalin for less than or equal to 14 day supplies | |
| | gabapentin pregabalin | | | |
| | TOPICAL LIDOCAINE | | | ZTLIDO |
| | Lidocaine Patches | | | |
| ADDITIONAL AGENTS | | Trial and failure of a tricyclic antidepressant greater than or equal to a 12 week supply AND trial and failure of gabapentin at a dose of 3600mg per day OR pregabalin for greater than or equal to a 12 week supply in the last 12 months will be required before approval can be given for a non-preferred agent. | carbamazepine imipramine (capsules) oxcarbazepine valproic acid | |
| | amitriptyline desipramine imipramine (tablets) nortriptyline | | | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|---|--|--|--|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| OPHTHALMICS | OP. -ANTI-ALLERGICS | | Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. Alomide and Alocril will be approved for pregnancy. Alomide will be approved for children under the age of 3. | ALOCRIL ALOMIDE bepotastine epinastine ZERVIAE |
| | ALREX azelastine BEPREVE* cromolyn 0.4% | | | |
| | OP. -ANTIBIOTICS- QUINOLONES | | Trial and failure of a preferred agent greater than or equal to 5 days in the last 12 months will be required before approval can be given for a non-preferred agent. | gatifloxacin ZYMAXID |
| | ciprofloxacin BESIVANCE gentamicin moxifloxacin 0.5% ofloxacin tobramycin | | | |
| | OP. -ANTI-INFLAMMATORY | | Trial and failure of ALL preferred agents each greater than or equal to 5 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | ACULAR/LS/PF (use preferred agent) ACUVAIL bromfenac 0.9% BROMSITE DUREZOL ILEVRO INVELTYS LOTEMAX SM loteprednol 0.5% (BRAND PREFERRED) PROLENSA |
| | flurbiprofen diclofenac LOTEMAX* ketorolac NEVANAC | | | |
| | OP. -BETA-BLOCKERS | | Trial and failure of three (3) preferred agents each greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. *Betoptic S will be approved for those with heart and lung conditions. | BETIMOL BETOPTIC S* |
| | betaxolol carteolol levobunolol timolol | | | |
| | OP. -CARBONIC ANHYDRASE INHIBITOR | | Trial and failure of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. | brinzolamide (BRAND PREFERRED) |
| | AZOPT dorzolamide | | | |
| | OP. -COMBO PRODUCTS | | | dorzolamide/timolol (BRAND PREFERRED) |
| | COMBIGAN* COSOPT* ROCKLATAN SIMBRINZA | | | |
| OP. -DRY EYE AGENTS | | Trial and failure of the preferred agent greater than or equal to 12 weeks will be required before approval can be given for the non-preferred agent. | CEQUA cyclosporine (BRAND PREFERRED) EYSUVIS MIEBO RESTASIS MULTIDOSE (use preferred agent) TYRVAYA | |
| RESTASIS* XIIDRA | | | | |
| OP. -PROSTAGLANDINS | | Trial and failure of ALL preferred agents each greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. | bimatoprost IYUZEH tafluprost | |
| latanoprost LUMIGAN TRAVATAN Z XALATAN ZIOPTAN | | | | |
| OP. -RHO KINASE INHIBITOR | | | | |
| RHOPRESSA | | | | |
| OP. -SYMPATHOMIMETICS | | Trial of a preferred agent greater than or equal to 30 days in the last 12 months will be required before approval can be given for a non-preferred agent. | brimonidine 0.15% (BRAND IS PREFERRED) | |
| ALPHAGAN P 0.1% ALPHAGAN P 0.15%* brimonidine 0.2% | | | | |
| OSTEOPOROSIS | BISPHOSPHONATES | | Trial and failure of a preferred agent greater than or equal to 12 months will be required before approval can be given for a non-preferred agent. Fosamax liquid will be approved for clients that have difficulty swallowing. **Evinity will only be allowed for a maximum of 12 months of treatment, will not be allowed with any concurrent osteoporosis treatment, and will be limited to approved indication ***Will be limited to 2 years of use | EVENITY** FORTEO*** FOSAMAX-D TYMLOS*** |
| | alendronate ibandronate risedronate | | | |
| | NASAL CALCITONIN | | | |
| calcitonin-salmon | | | | |
| OTIC | ANTIBIOTIC/STEROID COMBINATION | | ciprofloxacin 0.2% (use preferred agent) CIPRO HC (use preferred agent) CORTISPORIN-TC (use preferred agent) FLUOCINOLONE ACET OIL 0.01% (use preferred agent) | |
| | Ciprofloxacin/dexamethasone Neo/Poly/HC Suspension and Solution ofloxacin tobramycin/dexamethasone | | | |
| OVERACTIVE BLADDER | OVERACTIVE BLADDER AGENTS | | Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Oxytrol will be approved for clients that have an inability to swallow. | darifenacin GELNIQUE GEL 10% GEMTESA OXYTROL DIS tolterodine/ER trospium |
| | MYRBETRIQ oxybutynin /ER solifenacin TOVIAZ | | | |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|----------------------------|--|--|---|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERAL MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| PAIN | LONG-ACTING C-III | | <p>Trial and failure of a preferred agent(s) greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.</p> <p>C-III's and C-IV's that are not included on the PDL and are available without prior authorization with the exception of Butrans (generic substitution is mandatory).</p> <p>Concurrent therapy with a benzodiazepine and a narcotic medication or with duplicate benzodiazepines is not covered by Wyoming Medicaid. A single medication will continue to be allowed to process unless another benzodiazepine (or narcotic) is billed to Wyoming Medicaid.</p> <p>**Nucynta ER will be allowed for diabetic peripheral neuropathy or clients with significant gastrointestinal concerns with other CI narcotics.</p> <p>Fentanyl: 50mcg, 1 strength at a time, 1 patch every 3 days Hydromorphone ER: 30mg/day Hysingla ER: 120mg/day Methadone: Limited to 3 tablets per day Morphine ER: 90mg/day Nucynta ER: 327mg/day Oxycontin: 80mg/day Oxymorphone: 40mg/day Xtampza ER: 80mg/day</p> <p>Clients will be limited to one long-acting narcotic at a time</p> | <p>fentanyl patches hydrocodone ER hydromorphone ER HYSINGLA ER METHADONE morphine ER capsules (use preferred agents) NUCYNTE ER** oxymorphone ER OXYCONTIN XTAMPZA ER (additional criteria applies)</p> |
| | SHORT-ACTING C-III | | <p>Trial and failure of three (3) preferred agents greater than or equal to a 6 day supply in the last 90 days will be required before approval can be given for a non-preferred agent.</p> <p>*Nucynta will be allowed for diabetic peripheral neuropathy or clients with significant gastrointestinal concerns with other CI narcotics.</p> <p>Concurrent therapy with a benzodiazepine and a narcotic medication or with duplicate benzodiazepines is not covered by Wyoming Medicaid. A single medication will continue to be allowed to process unless another benzodiazepine (or narcotic) is billed to Wyoming Medicaid.</p> <p>All short-acting narcotics, after 42 days of consecutive use of any combination of short-acting narcotics, will be limited to 4 tablets per day (liquids have specific dosing limits per medication - please refer to dosage limitation chart at www.wymedicaid.org)</p> <p>Clients will be limited to one short-acting narcotic at a time</p> | <p>levorphanol NUCYNTE* oxymorphone ROXYBOND</p> |
| | C-III/C-V AGENTS | | <p>BUTRANS tramadol</p> | <p>Trial and failure of a preferred agent(s) greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent.</p> <p>Quantity and dosage limits apply (max 8 tabs/day).</p> <p>Concurrent therapy with a benzodiazepine and a narcotic medication or with duplicate benzodiazepines is not covered by Wyoming Medicaid. A single medication will continue to be allowed to process unless another benzodiazepine (or narcotic) is billed to Wyoming Medicaid.</p> |
| PARKINSON'S DISEASE | SHORT-ACTING AGENTS | | | |
| | LONG-ACTING AGENTS | | <p>**Non-preferred Parkinson's agents will require a 30 day trial and failure of 2 preferred medications including at least one short-acting agent and one long-acting agent</p> <p>*Neupro will be approved for clients with difficulty swallowing</p> | <p>APOKYN benztropine injectables GOCOVRI INBRIJA NEUPRO* ONGENTYS pramipexole ER XADAGO</p> |
| PHOSPHATE BINDERS | PHOSPHATE BINDERS | | <p>Prior authorization required for non-preferred agents.</p> | <p>AURYXIA lanthanum sevelamer VELPHORO</p> |

WYOMING MEDICAID
Preferred Drug List (PDL) February 7, 2024

| Please refer to the Additional Therapeutic Criteria Chart, Dosage Limitation List (red font indicates quantity/dosage limits apply), and the Wyoming Medicaid Provider Manual at http://wymedicaid.org for additional criteria. | | | | |
|--|---|---|--|--|
| THERAPEUTIC CLASS | PREFERRED AGENTS | PREFERRED AGENTS REQUIRING CLINICAL CRITERIA | CLINICAL CRITERIA | NON-PREFERRED AGENTS GENERIC MANDATORY POLICY APPLIES <small>THIS LIST IS NOT ALL INCLUSIVE PLEASE CONTACT CHANGE HEALTHCARE WITH ANY QUESTIONS</small> |
| PROSTATE | 5-ALPHA-REDUCTASE INHIBITORS | | Trial and failure of a preferred agent greater than or equal to a 30 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | dutasteride dutasteride/tamsulosin (<i>use separate agents</i>) |
| | finasteride | | | |
| | ALPHA BLOCKERS | | Trial and failure of a preferred agent greater than or equal to a 30 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. | alfuzosin dutasteride/tamsulosin (<i>use separate agents</i>) silodosin |
| | doxazosin tamsulosin terazosin | | | |
| PULMONARY ANTIHYPERTENSIVES | 5-ALPHA-REDUCTASE INHIBITORS | | Prior authorization required. Client must have a diagnosis of pulmonary hypertension. | sildenafil suspension (BRAND IS PREFERRED) |
| | | ALYQ tadalafil REVATIO SUSPENSION* sildenafil (Revatio A/B rated generic) | | |
| | ENDOTHELIN RECEPTOR ANTAGONISTS | | Prior authorization required. Client must have a diagnosis of pulmonary hypertension. | bosentan (BRAND IS PREFERRED) OPSUMIT (<i>use preferred agent</i>) TRACLEER TABS FOR ORAL SUSP (<i>use preferred agent</i>) |
| | | LETAIRIS TRACLEER TABS* | | |
| | GUANYLATE CYCLASE INHIBITORS | | Prior authorization required. | ADEMPAS (<i>use preferred agent</i>) |
| | PROSTACYCLINE VASODILATORS | | Prior authorization required. Client must have a diagnosis of pulmonary hypertension with documented right-heart catheterization validating the diagnosis. | |
| | ORENITRAM | | | |
| PROSTACYCLINE RECEPTOR AGONIST | | Prior authorization required. | UPTRAVI (<i>use preferred agent</i>) | |
| RESTLESS LEG SYNDROME | RESTLESS LEG SYNDROME | | Client must have a diagnosis of Restless Leg Syndrome (RLS). Trial and failure of gabapentin greater than or equal to 60 days and a trial and failure of a dopamine agonist greater than or equal to 60 days in the last 12 months will be required before approval can be given for a non-preferred agent. *Neupro will be approved for clients with difficulty swallowing or for clients with a diagnosis of Parkinson's Disease. Clients will not be allowed to take gabapentin and pregabalin concurrently | HORIZANT NEUPRO* |
| | pramipexole ropinirole | gabapentin | | |
| SKELETAL MUSCLE RELAXANTS | MUSCLE RELAXANTS | | Trial and failure of a preferred agent greater than or equal to a 14 day supply in the last 12 months will be required before approval can be given for a non-preferred agent. Cyclobenzaprine will require a prior authorization for clients concurrently taking a tricyclic antidepressant. Carisoprodol is limited to 84 tabs/365 days | carisoprodol chlorzoxazone cyclobenzaprine ER LYVISPAP metaxalone methocarbamol orphenadrine tizanidine capsules (<i>use preferred agent</i>) |
| | baclofen (tablets) cyclobenzaprine tizanidine tablets | | | |
| ULCERATIVE COLITIS | IMMUNOMODULATORS | | Client must have diagnosis of UC prior to approval of a preferred agent. To receive a non-preferred agent, client must have a diagnosis of UC and a 56-day trial and failure of the preferred agent. * Refer to Additional Therapeutics Clinical Criteria Chart for more information | ENTYVIO* REMICADE RINVOQ SIMPONI STELARA XELJANZ/XR |
| | | HUMIRA | | |
| UVEITIS | IMMUNOMODULATORS | | Client must have diagnosis of non-infectious intermediate, posterior, or panuveitis in adult patients | |
| | | HUMIRA | | |