Syphilis in Wyoming

Syphilis is a bacterial infection spread through condomless sex (oral, anal, and/or vaginal) and, less commonly, through injection drug use. The Wyoming Department of Health (WDH) recommends syphilis screening for all sexually active individuals. Also, WDH recommends repeated syphilis testing for pregnant women throughout pregnancy if the pregnant woman or partner(s) have other sexual partners, use injection or intranasal drugs, or have unprofessional tattoos or piercings

2022 Cases


During 2022, there were 66 cases of syphilis reported.

Female

Four cases were diagnosed during pregnancy in 2022; about 17% of cases in women

Almost half (48.8%) of male cases reported sex with male partners. Nationally, the rate of syphilis is **increasing** among men who have sex with men (MSM).


Male 65.2%




involved a pregnancy.

Diagnosis

Only **31.8%** of the cases were diagnosed with **primary** or **secondary syphilis.** Therefore, 68.2 % of cases were diagnosed after symptoms had resolved.





In 2022, the **25–34 age group** represented the largest portion of reported cases, followed by the **35–44 age group**.



Symptoms

Syphilis has **multiple stages of infection** that are defined by the presence or absence of specific symptoms. Common symptoms include painless sores (primary syphilis) and rash, hair loss, swollen lymph nodes, or mucous patches (secondary syphilis).





*The rates are per 100,000 population