Chlamydia in Wyoming


Chlamydia is a bacterial sexually transmitted infection (STI). Chlamydia is spread through condomless sex (oral, anal, and/or vaginal). It can also be spread from an infected mother to her baby during childbirth. Most people infected with chlamydia have no symptoms and do not seek testing. If symptoms do occur, they present within one to three weeks after exposure.

2023 Cases

During 2023, there were **1,829** cases of chlamydia reported.

In 2023, female cases comprised approximately **two-thirds** of Wyoming's reported chlamydia cases.

Female 65.8%




Male 34.2%

Chlamydia was the **most commonly reported** sexually transmitted infection (STI) in Wyoming in 2023.



Co-infections

Co-infections are the simultaneous presence of **two or more infections** in a patient. Approximately **three percent** of the chlamydia cases had a co-infection with gonorrhea.



In 2023, over **three-fifths** of chlamydia cases were reported in the **<25 age group**.

Repeat Infections

In 2023, over **five percent** of chlamydia cases had more than one infection with chlamydia during the calendar year.




Multiple Infections


Risk Factors

The data points associated with risk factors are **limited**.

Symptoms

Symptoms may include abnormal discharge from the infected site, burning during urination, itching, and pain during intercourse. If left untreated, chlamydia can cause **pelvic inflammatory disease (PID)**. Symptoms of PID only occur in patients assigned female at birth and include abdominal pain, fever, and chronic pelvic pain.





*The rates are per 100,000 population