

MMWR Week 7 (2/15/2026-2/21/2026)



INFECTIOUS DISEASE EPIDEMIOLOGY

Influenza Report

2025 - 2026 Influenza Season

MMWR Week 7 (2/15/26-2/21/26)

Updated February 27, 2026



Overview (MMWR Week 7)

Influenza and Influenza-like Illness (ILI) Activity

Spread

Regional

Transmission levels increased compared to last week

Flu Activity

Low

Reported influenza activity increased compared to last week

ILI Activity

Minimal

Reports of outpatient respiratory illnesses were minimal this week

Co-circulating

Other Respiratory Infections:

SARS-CoV-2, Pertussis, and increasing levels of RSV

Seasonal Data

Types of Flu:

Influenza A and B viruses are circulating; flu B percentages are increasing

Subtypes

Primary: A H3N2

Predominantly, H3N2 viruses were reported across the country this week

Outbreaks

0

No LTCF or school-associated influenza outbreaks were reported this week

Severity

Hospitalizations

The weekly influenza-associated hospitalization rate peaked in week 52 and has since decreased

Deaths

79

There have been 79 pediatric deaths reported across the country so far this season

Syndromic

1

There was one syndromic anomaly reported this week

EMS Reports

22

The number of suspected ILI reports increased compared to last week

Hot Spots

Tracking Trends

Most counties reported an increase in case counts compared to last week



Geographic Spread


Geographic Activity by Regions

Wyoming as a whole had **minimal ILI activity** this week (MMWR Week 7).
Transmission levels decreased compared to last week.

- Healthcare providers in **16** counties electronically reported influenza cases.
- The electronically reported influenza cases represent **all five** Infectious Disease Epidemiology (IDE) Geographic Regions.
- Healthcare providers across the state electronically reported **188** influenza cases (rapid influenza diagnostic tests and PCR-confirmed tests) this week.

IDE Geographic Regions of Wyoming

Southeast	Central	Western	Big Horn	North East
Regional, cases reported in Albany, Goshen, and Laramie	Local, cases reported in Fremont and Natrona	Regional, cases reported in every county	Local, cases reported in Park and Hot Springs	Regional, cases reported in Campbell, Weston, Sheridan, and Washakie



Virologic Surveillance

Public Health Laboratory


The overall volume of samples tested for influenza at the Wyoming Public Health Laboratory has steadily declined following the increasing distribution of the CDC Influenza SARS-CoV-2 Multiplex Assay, in addition to at-home testing options. We greatly encourage providers across the state to send additional specimens to the WPHL for virologic surveillance. The number of positive influenza specimens electronically reported this week **increased** compared to the previous week.

Healthcare and Clinical Laboratories

Clinical laboratories across the country most frequently reported **Influenza A/H3N2 viruses** circulating during MMWR Week 7, with increasing percentages of Influenza B viruses.

Electronic Lab Reports of Influenza Cases

Number of Electronic Lab Reported Cases



Influenza-Like Illness Surveillance

MMWR Week 7: 3.23% ▼ below WY baseline (5.36%)


The percent of patient visits to ILI-Net Sentinel Providers for ILI increased compared to last week.

The Wyoming Department of Health received reports from **more than 50%** of the ILINet providers across the state. Although weekly percentages could change as additional reports are submitted.

Key Updates: Seasonal influenza activity remains elevated nationally. Influenza A activity is decreasing while influenza B activity is increasing nationally and in most areas of the country. Based on CDC calculations, transmission within Wyoming was **minimal** this week. Nationally, the percentage of respiratory specimens testing positive for the influenza virus in clinical laboratories decreased slightly this week. Across HHS Regions, the overall percent of specimens testing positive for influenza increased in regions 1, 2, 3, 4, and 8; decreased in regions 6, 7, 9, and 10; and remained stable in region 5.

Weekly Percent of ILI Visits

Percent of Influenza-like Illness (ILI)



Pneumonia and Influenza Mortality


Mortality Data

Tracking death certificates is the most effective surveillance system for capturing and identifying pneumonia and influenza-associated deaths in Wyoming. According to the CDC, influenza is infrequently listed on death certificates. Also, testing for seasonal influenza infections is not frequently performed, particularly among the elderly, who are at greatest risk for seasonal influenza complications and death. Therefore, public health officials may not be able to identify influenza-associated deaths in many instances; as a result, this surveillance system may underestimate the true impact of influenza-associated deaths in the state.

There have been **44** pneumonia and influenza (P&I) mortality reports certified since the beginning of the 2025-2026 Influenza Season.

Monthly P&I Mortality Reports (2020-2026)





Number of P&I Deaths



Wastewater Surveillance

Here at the Wyoming Public Health Laboratory (WPHL), we have a team of scientists analyzing wastewater samples from across the state to identify trends in influenza (and many other pathogens). The graphs pictured below depict trend lines for the past two influenza seasons (2023-2026) as well as current trends. It is important to note that the high concentrations observed last season have dwarfed many of the previously detected values.

WPHL Sites: Cheyenne, Green River, Rawlins, and Sheridan



Wastewater Surveillance Continued

Our colleagues at the University of Wyoming (UW) have collaborated with the WPHL to analyze wastewater samples collected at four additional treatment facilities across the state. The graphs below show the current trend lines of varying influenza concentrations detected in wastewater, starting in June 2024 and continuing to the present. The current viral concentrations appear to be diminished compared to the higher values we observed last season during sustained community-wide transmission.

UW Sites: Laramie, Rock Springs, Jackson, and Gillette

